

REPORT DOCUMENTATION PAGE

Form Approved

OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)

2. REPORT DATE

11/5/90

3. REPORT TYPE AND DATES COVERED 10/1/89-
Technical Rept. (Interim) 9/30/90

4. TITLE AND SUBTITLE

Complex Route to Chaos in Velocity Driven Atoms

5. FUNDING NUMBERS

C: N00014-89-WX-24146
N00014-90-WX-24103

PE: 0601153N

PR: RR011-03-OK

TA: 412n006---02

WU: 61-3299-00

6. AUTHOR(S)

Phouc X. Tran, D.W. Brenner and C.T. White

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Naval Research Laboratory
Washington, DC 20375-5000
(Code 6119, C.T. White)8. PERFORMING ORGANIZATION
REPORT NUMBER

Technical Report #12

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)

Office of Naval Research
800 North Quincy Street
Arlington, VA 22217-5000
(Code 1112AI, D.H. Liebenberg)10. SPONSORING / MONITORING
AGENCY REPORT NUMBER

Technical Report # 12

11. SUPPLEMENTARY NOTES

12a. DISTRIBUTION / AVAILABILITY STATEMENT

This document has been approved for public release and
sale, distribution of this document is unlimited.

12b. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 words)

The oscillations of a linear triatomic molecule with the first atom driven at a constant velocity v are studied numerically. As v increases, we observe a sequence of transitions from quasiperiodicity to chaos, mode locking, chaos, model locking, chaos, another mode locking then to chaos before the atoms break off. The largest Lyapunov exponent does not rise linearly with the driven velocity. The mode locking structure for the last onset to chaos has a universal fractal dimension 0.871 ± 0.001 at a winding number far from the golden mean.

14. SUBJECT TERMS

molecular dynamics, chaos, triatomic molecule

15. NUMBER

16. PRICE CODE

17. SECURITY CLASSIFICATION
OF REPORT

UNCLASSIFIED

18. SECURITY CLASSIFICATION
OF THIS PAGE

UNCLASSIFIED

19. SECURITY CLASSIFICATION
OF ABSTRACT

UNCLASSIFIED

20. LIMITATION OF ABSTRACT

DTIC
ELECTE
DEC 13 1990
S E D

OFFICE OF NAVAL RESEARCH

CONTRACT #'s:

N00014-89-WX-24146

N00014-90-WX-24103

R&T CODE 412n006---02

TECHNICAL REPORT # 12

COMPLEX ROUTE TO CHAOS IN VELOCITY DRIVEN ATOMS*

Phouc X. Tran, D.W. Brenner, and C.T. White

Code 6119

Naval Research Laboratory
Washington, D.C. 20375-5000

Prepared for publication



Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input checked="checked" type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special

A-1

*Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

Complex route to chaos in velocity-driven atoms*

Phuoc X. Tran, D.W. Brenner and C. T. White

Naval Research Laboratory

Washington, DC. 20375-5000

August 1990

Abstract

The oscillations of a linear triatomic molecule with the first atom driven at a constant velocity v are studied numerically. As v increases, we observe a sequence of transitions from quasiperiodicity to chaos, mode locking, chaos, another mode locking then to chaos before the atoms break off. The largest Lyapunov exponent does not rise linearly with the driven velocity. The mode locking structure for the last onset to chaos has a universal fractal dimension 0.871 ± 0.001 at a winding number far from the golden mean.

*This work was supported by ONR through NRL and the ONR Physics Division.

PACS Numbers: 05.45.+b, 82.40.Fp

Many dissipative systems are known to undergo transitions to chaos by different routes. Such transitions in a system with few degrees of freedom may provide some insight into turbulence. The quasiperiodic transition has been observed experimentally^{1,2} in the Rayleigh-Bénard convective fluid, thermo-acoustic oscillation, electronic conduction in extrinsic Ge, Josephson junctions, charge density waves, and many others. These are some of the experiments that could be modeled by the circle map. Recent theoretical developments have resulted in some quantitative and universal characterizations for the onset of chaos³. In most of experiments on the quasiperiodic transition, one of the two frequencies can be varied directly. This allows the ratio of the two frequencies (or winding number) to be set arbitrarily close to a value corresponding to the golden mean ($0.61803\dots$) where one can expect to observe a universal scaling behavior at the onset of chaos as predicted by theory. In some experiments, however, the two frequencies arise intrinsically from the system and can not be controlled directly. In these cases, for example the Rayleigh-Bénard convective ⁴He fluid⁴, it is somewhat difficult to achieve the golden mean ratio. However, the system exhibits a quasiperiodic transition to chaos with a mode locking staircase that has a fractal dimension D close to 0.868 in good agreement with theoretical predictions by the circle map. As predicted, this universal value of D is a global property and should hold not only at the golden mean but also for a range of winding number at the onset of chaos^{3,5}. Furthermore, many condensed matter systems are known to undergo transitions to chaos when there is a competition between two incommensurate frequencies which is not necessarily at the golden mean.

In our model, we observed a different route to chaos beginning with two incommensurate frequencies. The external control parameter can only vary both frequencies indirectly. After the first transition to chaos, there is an abrupt transition to mode locking that gives rise to the first section of an incomplete devil's staircase amidst a set of quasiperiodic points. This is followed by a window of chaos then another section of a mode-locked staircase, then comes the last stage of chaos before the disappearance of the attractor signaling a separation of the atoms.

The atomic motions in a gas or fluid driven by a high pressure front are likely to become turbulent. Molecular dynamics studies of shock front propagation in solids often involve the so called piston problem. In the one-dimensional version of this problem, one atom at the end of a long chain is driven at a constant velocity and the dynamics of the remaining atoms in the system is studied. The problem attracting our interest is a simpler but related one. It consists of three identical atoms moving in a straight line with the first atom driven at a constant velocity v . These atoms are assumed to interact through a potential⁶

$$V_{ij} = C[1 - K(x_i - x_j)]e^{-K(x_i - x_j)} \quad (1)$$

where x_i is the position of the i^{th} atom, K and C are positive constants proportional to the equilibrium separation and the binding energy respectively. As in the model containing many more atoms, this is a non-conservative system because a varied force is required to maintain a constant velocity of the first atom. Here we will investigate in detail the relative motions of the three atoms, focussing on the behavior of the bond length in response to different values of v . An immediate question concerns the value of v that causes the three atoms to break apart, and the chaotic behavior that may occur in the system.

We begin by integrating the following equations of motion

$$\frac{d^2 Y_i}{dT^2} = (Y_{i+1} - Y_i - 2)e^{-(Y_{i+1} - Y_i)} - (Y_i - Y_{i-1} - 2)e^{-(Y_i - Y_{i-1})} \quad (2)$$

where $Y_i = Kx_i$ and $T = \omega t$ are the new atomic position and time in dimensionless units respectively. The dimensions used in this calculation can be scaled so that each unit of velocity, distance and time correspond to 8.295 Km/sec, 0.55 Å and $\omega^{-1} = 6.63 \times 10^{-15}$ sec respectively. These values were based on an Nitrogen atomic mass of 14 amu, a bond length of 1.1 Å and binding energy of 5 eV. Any other type of atom will simply result in a different set of scaled units. We will consider the oscillations of the two bond lengths $B_1 = Y_2 - Y_1$ and $B_2 = Y_3 - Y_2$. In solving the set of ordinary differential equations, one may not wish to use the relative coordinates due to a constraint on the velocity of the first atom.

For an extremely small value of v , the atoms are near their equilibrium positions and their interaction can be approximated by a harmonic potential. Then the ratio of two natural frequencies of the system is known as $1/\sqrt{3}$. The two bonds start oscillating with two incommensurate frequencies. We will limit our discussion to the first bond near the shock front. As we increase v , harmonic frequencies of the form $mf_1 + nf_2$ begin to appear as seen in the power spectrum in Fig.1a. The phase space diagram shows a quasiperiodic attractor for the bond which oscillates about its equilibrium value of 2.0 (Fig.2a). The Poincaré section of this attractor is shown in Fig.3a. The first transition to chaos occurs when v reaches 0.267. The power spectrum develops broad band signals (Fig.1b), corrugations of the Poincaré section appear (Fig.3b) and trajectories in the phase space start wandering (Fig.2b). Fully developed chaos takes place when v reaches 0.29 with characteristics shown in Fig.1c, 2c and 3c.

A sudden transition to quasiperiodicity takes places when v reaches 0.3093. Then a mode locking begins at $v = 0.3098$ and continues until $v = 0.31281$. The corresponding winding numbers are $7/18$ and $5/13$ respectively. We found two other mode-locked ratios within this interval using the Farey tree sequence. The prospect of locating other mode-locked ratios within this interval is certain, but very time consuming. Interspersed among the mode-locked ratios are quasiperiodic points (Fig.4).

The second transition to chaos takes place when v is slightly greater than 0.31281. Similar chaotic behaviors in the phase space, Poincaré map and power spectrum are obtained. A second series of mode-locked states emerges when $0.31309 < v < 0.31700$. In this interval we also show four mode-locked ratios using the Farey tree. Of course, there are other mode-locked states and quasiperiodic states as before. For each mode-locked ratio, The Poincaré maps and phase space diagrams clearly show the number of resonances of trajectories corresponding to the numerators of the mode-locked ratios (Fig.5). The existence of quasiperiodic points among the plateaux in the two mode-locked intervals suggests that the devil's staircase is incomplete and the transition to chaos is far from criticality.

The third transition to chaos takes place as v is increased past 0.317. The chaotic states persist until $v = 0.36804$. By increasing v from below to above this value, the chaotic attractor severely deforms. Its amplitude of oscillation bursts into a linear motion as a result of a separation of the first two atoms. In another words the attractor disappears. The bond stops oscillating about its equilibrium position and keeps stretching out as time progresses.

Since the two portions of the mode-locked staircase are separated by an interval of chaos in which the fractal characteristic is discontinued, we compute the fractal dimension of each of the two sections of the staircase separately using the expression $\sum_{i=1}^n (s_i/s_t)^D = 1$, where s_i is the gap between two plateaux and s_t is the largest gap. We obtained $D = 0.915 \pm 0.001$ for the first section and $D = 0.871 \pm 0.001$ for the second one. The value of D for the second portion of the staircase is in good agreement with the universal value $D = 0.868 \pm 0.002$. The larger value of D for the first section may arise from a transition far from criticality, as this same value of D has been found in the scaling behavior for the charge density wave and Josephson junction². It appears that the nonlinearity in this model is not a monotonic function of the control parameter v . Then it is not feasible to show whether the system is approaching chaos from below or above criticality, although D is less than unity.

The sequence of transitions is summarized in Fig.6 with three chaotic regimes CH1 to CH3 alternating among a quasiperiodic QP2 regime and two mode-locked staircases PL1 and PL2. The region to the right of CH3 marks a disappearance of the attractor, as the atoms become separated. We calculate the largest Lyapunov exponent λ for the attractors within each of the three chaotic intervals using the method based on the evolution of trajectories⁷. As shown in Fig.7, there is a nonlinear relation between the control parameter v and the level of chaos. After some fluctuations during the first chaotic interval, λ decreases in the second and last interval while remaining positive. In this model, the nonlinearity parameter is fixed (K), the factor that leads to a transition to chaos is due to the interaction of the external driven force, parameterized by the

velocity v , with a nonlinear potential. Significant nonlinearity will arise when the driven force with a large value of v interacts with higher order terms of K from the exponent in the potential. Driven nonlinear systems often generate both subharmonics and higher harmonics. In the first transition to chaos, higher harmonics ($mf_1 + nf_2$) were gradually suppressed. The lower fundamental frequency f_1 was the most stable component to disappear at the onset of chaos. We do not know the exact mechanism for the repeated transition to chaos or for the reappearance of higher harmonics when the system goes from chaos to mode-locking. However, the indirect cause may be linked to the oscillation of the nonlinearity generated under different values of the driven velocity. This is reflected in the curve for the Lyapunov exponent.

There are systems in which chaos emerges regularly in parameter space. A well known example is the Belousov-Zhabotinsky reaction. This experiment⁸ reveals that alternating regimes of chaos and periodicity appear regularly in a sequence of multiple bifurcations. In this case, the transition to chaos may be characterized by a different universal number but does not possess a mode locking structure and a fractal dimension as in our model. It is also known that experiments on the transition to chaos via mode locking exists^{2,9} but the recurrence of chaos was not found in these models. In a particular Rayleigh-Bénard experiment¹⁰ for a specific type of convective cells, it has been observed that the system transitioned from quasiperiodicity to chaos then to mode-locking at one winding number only, but no mode-locked structure has been found. We know of no previous results from experiments or theoretical models that exhibit a similar sequence of transitions to chaos like our model. We have also looked into a similar model with a much larger size, up to five hundred atoms in a linear chain. Similar attractors were found on groups of three atoms when the chain was driven by a strong front and broken into segments. However, the dynamics is very complex and the work is still underway. For a system with few degrees of freedom, this is the first model that exhibits recurrent transitions to chaos and a mode-locked staircase with a universal fractal dimension. It also provides some new details to the dynamical process of atoms interacting with a steady moving front.

Acknowledgement:

This work was supported by ONR through the ONR physics division, contract #N00014-89-WX-24146.

References

- [1] J. Stavans, F. Heslat, and A. Lichaber, *Phys. Rev. Lett.* **55**, 596 (1985); T. Yazaki, S. Sugioka, F. Mizutani, and H. Mamada, *Ibid.* **64**, 2515 (1990); E.G. Gwinn and R.M. Westervelt, *Ibid.* **57**, 1060 (1986); D.J. Olinger and K.R. Sreenivasan, *Ibid.* **60**, 797 (1988).
- [2] S.E. Brown, G. Mozurkewich, and G. Grüner. *Ibid.* **52**, 2277 (1984); W.J. Yeh, Da-Ren He, and Y.H. Kao, *Ibid.* **52**, 480 (1984).
- [3] P. Cvitanović, M. Jensen, L. Kadanoff, and I. Procaccia, *Ibid.* **55**, 343, (1985); M. Jensen, P. Bak, and T. Bohr, *Phys. Rev. A* **30**, 1960 (1984); T. Halsey, P. Meakin, and I. Procaccia, *Phys. Rev. Lett.* **56**, 854 (1986).
- [4] H Haucke and R. Ecke, *Physica.* **25D**, 307 (1987).
- [5] H.L. Swinney and J. Maselko, *Phys. Rev. Lett.* **55**, 2366 (1985).
- [6] P.X. Tran, D.W. Brenner and C.T. White, *Int. J. Quant. Chem.*, Proceedings of the Sanibel symposia, in press (1990).
- [7] G. Benettin, L. Galgani and J.M Strelcyn, *Phys. Rev. A* **14**, 2338 (1976).
- [8] C. Vidal, S. Bachelart and A. Rossi, *J. Phys. (Paris)* **43**, 7 (1982).
- [9] J.P. Gollub and S.V. Benson, *J. Fluid Mech.* **100**, 449, (1980).
- [10] P. Bergé, Y. Pomeau and C. Vidal, *Order within chaos*, J. Wiley and Hermann (Paris), p.180 (1984).

List of Figures:

Figure 1a: Power spectrum for the oscillating bond: a) Quasiperiodic regime, here $v = 0.23$. b) The first transition to chaos, $v = 0.267$. c) Chaotic state, $v = 0.28$. These same values of v are used in Figs. 2 and 3.

Figure 2: Two-dimensional phase portraits from quasiperiodic a) to chaotic state c).

Figure 3: Poincaré sections obtained from cutting the attractors in Fig.2.

Figure 4: Plot of the winding number $w = \frac{f_1}{f_2}$ for the two sections of the devil's staircase. We only show the large plateaux and a small number of quasiperiodic points. Between the two sections is a chaotic region.

Figure 5: A typical mode-locked state is shown here for $w = 3/8$, a) Phase space portrait, b) Poincaré section. Note how the attractor transforms from quasiperiodicity (Fig.2a) to mode-locking (Fig.5a).

Figure 6: Summary of the transition sequence plotted on the velocity v axis. See text for details.

Figure 7: Variation of the largest Lyapunov exponent λ versus the velocity v . In the two regimes of mode-locked states, λ drops to near zero.




Fig.1-3




Fig.4




Fig.5




Fig.6




Fig.7

REPORTS DISTRIBUTION FOR ONR PHYSICS DIVISION
UNCLASSIFIED CONTRACTS

Director Defense Advanced Research Projects Agency Attn: Technical Library, T10 1400 Wilson Blvd. Arlington Virginia 22209-2309	1 copy
Office of Naval Research Physics Division Office (Code 1112) 800 North Quincy Street Arlington, Virginia 22217-5000	2 copies
Office of Naval Research Director, Technology (Code 20) 800 North Quincy Street Arlington, Virginia 22217-5000	1 copy
Naval Research Laboratory Department of the Navy (Code 2625) Attn: Technical Library Washington, D.C. 20375-5000	1 copy
Office of the Director of Defense Research and Engineering Information Office Library Branch The Pentagon, Rm. 3E 1006 Washington, D.C. 20301	1 copy
U.S. Army Research Office Box 12211 Research Triangle Park North Carolina 27709-2211	2 copies
Defense Technical Information Center Cameron Station Alexandria, Virginia 22314	4 2 copies
Director National Bureau of Standards Research Information Center Attn: Technical Library (Admin E-01) Gaithersburg, MD 20899	1 copy
Commander U.S. Army Belvoir Research, Development and Engineering Ctr. Attn: Technical Library (STRBE-BT) Fort Belvoir, Virginia 22060-5606	1 copy

ODDR&E Advisory Group on Electron Devices
201 Varick Street, 11th Floor
New York, New York 10014-4877

1 copy

Air Force Office of Scientific Research
Department of the Air Force
Bolling AFB, DC 22209

1 copy

Air Force Weapons Laboratory
Technical Library
Kirtland Air Force Base
Albuquerque, NM 87117

1 copy

Lawrence Livermore Laboratory
Attn: Dr. W. F. Krupke
University of California
P.O. 808
Livermore, CA 94550

1 copy

Harry Diamond Laboratories
Technical Library
2800 Powder Mill Road
Adelphi, MD 20783

1 copy

Naval Weapons Center
Technical Library (Code 753)
China Lake, CA 93555

1 copy

Naval Underwater Systems Center
Technical Center
New London, CT 06320

1 copy

Commandant of the Marine Corps
Scientific Advisor (Code RD-1)
Washington, DC 20380

1 copy

Naval Ordnance Station
Technical Library
Indian Head, MD 20640

1 copy

Naval Postgraduate School
Technical Library (Code 0212)
Monterey, CA 93940

1 copy

Naval Missile Center
Technical Library (Code 5632.2)
Point Mugo, CA 93010

1 copy

Naval Ordnance Station
Technical Library
Louisville, KY 40214

1 copy

Commanding Office
Naval Ocean Research & Development Activity
Technical Library
NSTL Station, MS 39529

1 copy

Naval Oceans Systems Center
Technical Library
San Diego, CA 92152

1 copy

Naval Surface Weapons Center
Technical Library
Silver Spring, MD 20910

1 copy

Naval Ship Research and Development Center
Central Library (Codes L42 and L43)
Bethesda, MD 20084

1 copy

Naval Avionics Facility
Technical Library
Indianapolis, IN 46218

1 copy

a